**CLAIMS:**

1. (Original) Compounds of the general formula (I):




(I),

wherein

$R^1$ is a hydrogen atom, a heteroalkyl, heteroalkylcycloalkyl or heteroaralkyl radical,

the radicals $R^2$, each independently of any other(s), are halogen atoms, hydroxy, amino, nitro or thiol groups, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radicals,

the radicals $R^3$, each independently of any other(s), are halogen atoms, hydroxy, amino, nitro or thiol groups, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radicals,

G is a glycosyl group,

n is 0, 1, 2, 3 or 4 and

m is 0, 1, 2, 3 or 4,

or a pharmacologically acceptable salt, solvate, hydrate or pharmacologically acceptable formulation thereof.

2. (Currently amended) Compounds according to claim 1, ~~wherein~~ wherein $R^1$ is a hydrogen atom or a group of formula $COOR^4$ or $CONR^5R^6$ wherein $R^4$, $R^5$ and $R^6$ are, each independently of the others, hydrogen atoms, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radicals, or $R^5$ and $R^6$ together are part of an optionally substituted heteroaryl or heterocycloalkyl ring.

3. (Currently amended) Compounds according to claim 1 ~~or 2~~, wherein $R^4$ is a hydrogen atom, a $C_1$-$C_4$alkyl or benzyl radical.

4. (Currently amended) Compounds according to ~~one of claims~~ claim 1 ~~to 3~~, wherein $R^1$ is a hydrogen atom or a group of formula COOH or COOEt.

5. (Currently amended) Compounds according to ~~one of claims~~ claim 1 ~~to 3~~, wherein $R^1$ is a group of formula $CONHR^5$ and wherein $R^4$, $R^5$ and $R^6$ are, each independently of the others, hydrogen atoms, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radicals, or $R^5$ and $R^6$ together are part of an optionally substituted heteroaryl or

heterocycloalkyl ring ~~$R^5$ is defined as in one of the preceding claims~~.

6. (Currently amended) Compounds according ~~to one of claims~~ claim 1 ~~to 5~~, wherein m is 0.

7. (Currently amended) Compounds according to ~~one of claims~~ claim 1 ~~to 5~~, wherein m is 1 and $R^3$ is a hydroxy group which is bonded to the phenyl ring in a position ortho to the amidino group.

8. (Currently amended) Compounds according to ~~one of claims~~ claim 1 ~~to 7~~, wherein n is 2.

9. (Currently amended) Compounds according to ~~one of claims~~ claim 1 ~~to 8~~, wherein the radicals $R^2$, each independently of any other(s), are $C_1$-$C_4$alkyloxy, $C_1$-$C_4$hydroxyalkyloxy or benzyloxy groups.

10. (Currently amended) Pharmaceutical compositions comprising a compound according to ~~claims~~ claim 1 ~~to 9~~ as active ingredient and, optionally, carrier substances and/or adjuvants.

11. (Currently amended) A method for inhibiting blood clotting in a patient in need thereof comprising administration of an effective amount of a pharmaceutical composition as claimed in claim 10 ~~Use of a compound or of a pharmaceutical composition according to one of claims 1 to 10 in inhibiting factor VIIa~~.

12. (Currently amended) A method as claimed in claim 11, wherein said composition is administered ~~Use of a~~

~~compound or of a pharmaceutical composition according to one of claims 1 to 10 in the preparation of a medicament~~ for the treatment and/or prevention of a thromboembolic conditions~~,~~ selected from the group consisting of arterial restenosis, septicaemia, cancer, and acute inflammation ~~or other conditions mediated by factor VIIa activity~~.

13. (Currently amended) The method as claimed in claim 11, wherein said composition is administered to a patient undergoing ~~Use of a compound or of a pharmaceutical composition according to one of claims 1 to 10 in the preparation of a medicament for utilisation in~~ vascular surgery.